

Dynamical mean-field theory of Hubbard-Holstein model at half-filling: Zero temperature metal-insulator and insulator-insulator transitions

Gun Sang Jeon,^{1,2} Tae-Ho Park,¹ Jung Hoon Han,^{1,3} Hyun C. Lee,⁴ and Han-Yong Choi^{1,5,6}

¹*Department of Physics, BK21 Physics Research Division,
and Institute for Basic Science Research, Sung Kyun Kwan University, Suwon 440-746, Korea.*

²*The Pennsylvania State University, Department of Physics, University Park, PA 16802.*

³*CSCMR, Seoul National University, Seoul 151-747, Korea.*

⁴*Department of Physics and Basic Science Research Institute, Sogang University, Seoul 121-742, Korea*

⁵*CNNC, Sung Kyun Kwan University, Suwon 440-746, Korea.*

⁶*Asia Pacific Center for Theoretical Physics, Pohang 790-784, Korea.*

(Dated: February 2, 2008)

We study the Hubbard-Holstein model, which includes both the electron-electron and electron-phonon interactions characterized by U and g , respectively, employing the dynamical mean-field theory combined with Wilson's numerical renormalization group technique. A zero temperature phase diagram of metal-insulator and insulator-insulator transitions at half-filling is mapped out which exhibits the interplay between U and g . As U (g) is increased, a metal to Mott-Hubbard insulator (bipolaron insulator) transition occurs, and the two insulating states are distinct and can not be adiabatically connected. The nature of and transitions between the three states are discussed.

The interaction-induced phenomena are a fundamental problem because of the new physics that emerges. For instance, the interaction-driven metal-insulator transition (MIT) has been studied to a great extent for systems which involve the electron-electron or electron-phonon interactions [1]. In real condensed-matter systems, both interactions exist and it will be important to understand the interplay between them. The prototype model for describing such interplay is the Hubbard-Holstein (HH) model, which includes the onsite electron-electron and electron-phonon interactions characterized by U and g , respectively, as given by Eq. (1) below. The Hubbard model ($g = 0$) and Holstein model ($U = 0$), which are special limits of the more general HH model, have been extensively studied using the dynamical mean-field theory (DMFT) in infinite dimensions [2, 3] in the context of the interaction-driven MIT. Both models exhibit MIT as U or g is increased above a critical value. Now, consider the MIT in the $U - g$ plane. Natural questions then arise as to whether and how the two insulating states, metallic states, and metal-insulator transitions of the Hubbard and Holstein models are different, and how they are affected when both U and g are present and compete with each other. These questions will be addressed here by studying the HH model within the DMFT combined with Wilson's numerical renormalization group (NRG) [4]. The NRG technique [5, 6] is particularly powerful in that it is non-perturbative in nature so that it can cover the whole parameter space and that it can probe the extremely small energy scales like the narrow coherence peak and soft phonon mode which emerge as the metal-insulator transitions are approached.

We will present a zero temperature phase diagram of unbroken symmetry ground states with a focus on the nature of and transitions between the ground states as U and g are varied. The main results of the present work

are: (i) The ground state is a metallic (M) state when both U and g are small, but is a bipolaron insulating (BPI) state when g is large, and is a Mott-Hubbard insulating (MHI) state when U is large (see Fig. 1). In the shaded region in Fig. 1, the insulating and metallic solutions coexist, and the ground state is the one with a lower energy. (ii) The phase transition between M and BPI occurs along the dashed line that cuts through the coexistence region connecting E and T of Fig. 1. The order parameter ($\langle n - 1 \rangle$, see below.) changes discontinuously at the transition and the M-BPI transition is first order (Fig. 2). (iii) The transition between M and MHI occurs along the outer boundary of the coexistence region denoted by the solid line connecting T and U2. It is second order as in the Hubbard model (Fig. 3). (iv) The two kinds of insulators (BPI and MHI) are distinct and can *not* be adiabatically connected. The transition between them is first order, and occurs along $U = 2g^2/\omega_0$ (Fig. 4. See below). These points will be discussed in more detail below.

The HH model is defined by

$$\mathcal{H} = -\frac{t}{\sqrt{q}} \sum_{\langle i,j \rangle \sigma} c_{i\sigma}^\dagger c_{j\sigma} + U \sum_i n_{i\uparrow} n_{i\downarrow} + \omega_0 \sum_i a_i^\dagger a_i + g \sum_i (a_i^\dagger + a_i) (n_i - 1), \quad (1)$$

where q is the nearest-neighbor coordination number, $n_{i\sigma} = c_{i\sigma}^\dagger c_{i\sigma}$ the electron density operator for spin σ at site i , and $n_i = \sum_\sigma n_{i\sigma}$. The electrons have the onsite Coulomb repulsion U , and are linearly coupled with the Einstein phonon of frequency ω_0 with the onsite coupling constant g . The DMFT was employed to solve the HH model of Eq. (1). The DMFT maps a lattice model onto an effective single site impurity model imbedded in a bath which is determined self-consistently via iterations [2, 3].




FIG. 1: The zero-temperature metal-insulator transition phase diagram of the HH model at half-filling. The cuts labeled with 2 (a & b), 3, and 4 correspond, respectively, to the Figs. 2, 3, and 4. The cuts 2a, 2b, 3, and 4 represent, respectively, the calculations along the fixed $g = 0.16$, $U = 1.0$, $g = 0.1$, and $U = 2.5$. The shaded area is the coexistence region where both the metallic and insulating solutions exist. The dashed lines between M and BPI, and between BPI and MHI states represent 1st order phase transitions, and the solid line between M and MHI represents 2nd order. See the text for details.

The effective impurity problem was solved by the NRG technique in the present study.

The NRG+DMFT has been successfully applied to the half-filled Hubbard model to study the MIT at both $T = 0$ [4] and finite temperatures [7], and to the half-filled Holstein model at $T = 0$ [8]. For the Hubbard model, the ground state is insulating for $U > U_{c2}$ and metallic for $U < U_{c1}$. Between $U_{c1} < U < U_{c2}$, the metallic and insulating solutions coexist, and the metallic state has the lower energy and is the ground state. The MIT occurs at $U_c = U_{c2}$, and is 2nd order. For the Holstein model, the ground state is insulating for $g > g_{c2}$ and metallic for $g < g_{c1}$. Meyer *et al.* reported that $g_{c2} - g_{c1}$ is reduced as the phonon frequency ω_0 is decreased and $g_{c1} = g_{c2}$ for $\omega_0 = 0.05 W$, where $W = 4t$ is the bandwidth [8]. We generalized these lines of research and study the HH model at half-filling with the NRG+DMFT incorporating the improved method for calculating electron [9] and phonon spectral function [10]. We adopt the semi-circular density of states which is realized in the Bethe lattice in infinite dimensions.

Our central result is the Fig. 1, where we present a zero temperature phase diagram in the $U - E_{bp}$ plane of unbroken symmetry ground states of the HH model at half-filling obtained from the NRG+DMFT calculations. $E_{bp} = 2g^2/\omega_0$ is the bipolaron binding energy which is twice the polaron energy, $E_p = 2E_{bp}$. We take the bandwidth W as the unit of energy ($W = 1$). We

take $\omega_0 = 0.05$ as with Meyer *et al.* Three distinct states are found in the $U - E_{bp}$ plane: metal, bipolaron insulator, and Mott-Hubbard insulator states. We will look at them in more detail below. Specifically, we will address (i) the nature of M, BPI, and MHI states, (ii) transition between M and BPI, (iii) transition between M and MHI, and (iv) transition between the BPI and MHI. The results obtained from the NRG+DMFT regarding these points were summarized in the introduction above.

(i) *Nature of M, BPI, and MHI states.* – The ground state is a metallic state when both U and g are small. The M state becomes more correlated as g or U is increased as reflected in the decreasing quasiparticle weight $z = [1 - \partial \Sigma_1(\omega)/\partial \omega]_{\omega \rightarrow 0}^{-1}$, where the subscript 1 (2) refers to the real (imaginary) part. z vanishes at the outer boundary of the coexistence region. The BPI state is insulating because the attractive interaction between the electrons mediated by the phonons binds two electrons into a bipolaron as Capone and Ciuchi reported [11]. The MHI state, on the other hand, is insulating because of the repulsion U between two electrons at the same site.

An important point for understanding the overall feature of the phase diagram is that the effective interaction between two electrons in the HH model is given, after integrating the phonons out, by

$$V_{eff}(\omega) = U + \frac{2g^2\omega_0}{\omega^2 - \omega_0^2}. \quad (2)$$

The interactions between electrons due to g and U compete each other for $\omega \lesssim \omega_0$, which yields $V_{eff}(\omega = 0) = 0$ along $U = E_{bp}$. The $U = E_{bp}$ line plays a special role in the phase diagram in that all three states meet at the triple point T lying on this line and the BPI-MHI transition occurs along it, as we will discuss below. Also, the critical g_c , which separates M and BPI along E and T, should increase as U is increased because the attractive interaction due to g has to overcome the repulsive interaction of U to bind two electrons into a bipolaron. This expectation is indeed borne out as shown in Fig. 1. On the other hand, the U_c , which is the solid line connecting T and U2, depends weakly on g .

For the coexistence region with $U > E_{bp}$, the arguments which led to the conclusion of the MIT of the Hubbard model being a 2nd order at $U = U_{c2}$ [12] still apply and the M is the ground state. On the other hand, in the coexistence region with $E_{bp} > U$, the $\langle n_{\uparrow}n_{\downarrow} \rangle_{BPI}$ and $\langle n_{\uparrow}n_{\downarrow} \rangle_M$, where $\langle \mathcal{O} \rangle_{M(I)}$ stands for an expectation value of an operator $\mathcal{O}$ in the metallic (insulating) state, do not meet as shown in Fig. 2(a). This implies that the M-BPI transition is 1st order. The energy calculations in Fig. 2(a) show that the M and BPI solutions cross each other inside the coexistence region.

(ii) *Transition between M and BPI.* – The MIT of half-filled Holstein model in infinite dimensions was discussed




FIG. 2: (a) The $\langle (n-1)^2 \rangle / 2$, which is equal to the double occupancy $\langle n_\uparrow n_\downarrow \rangle$ in M state, and ground state energy E_g are plotted along the cut 2a with a fixed $g = 0.16$ in Fig. 1 as U is increased (for M) or decreased (for BPI). It can be read off from the plots that the coexistence region is $0.925 < U < 1.0$, and the phase transition occurs around $U = 0.96$ where the energies of the M and BPI states become the same. (b) Plots of z , C , and $\frac{\omega_0}{g}\langle a \rangle$ as g is increased along the cut 2b of Fig. 1. In Figs. (c) and (d), the electron and phonon spectral functions, $A(\omega)$ and $B(\omega)$, respectively, are shown along the same cut with (b).

previously by Meyer *et al.* [8] and by Benedetti and Zeyher [13]. The BPI state is insulating because electrons are bound in bipolarons. Because we set the chemical potential $\mu = U/2$, which prefers the electron density $\langle n \rangle = 1$, the bipolaron formation may be accommodated by reconstructing the system into a phase separated state or a charge ordered state where the doubly occupied and empty sites alternate in the real space. This possibility is not allowed in the present DMFT work because the unit cell consists of a single site. Instead, the bipolaron instability shows up as degenerate ground states: In the NRG+DMFT calculations, the ground state is 8 fold degenerate in the BPI regime, with a one-to-one correspondence between the two sets of $\langle n \rangle - 1 \approx 1$ and $\langle n \rangle - 1 \approx -1$, which on averaging satisfies $\langle n \rangle = 1$. The order parameters for the M-BPI may be taken as

$$C = \langle n - 1 \rangle. \quad (3)$$

Now, consider the cut 2b in Fig. 1 along which we increase g with $U = 1.0$. Let g_{c1} and g_{c2} be, respectively, the lower and upper boundaries of the shaded coexistence region. As $g \rightarrow g_{c2}$, the quasiparticle weight z and the renormalized phonon frequency Ω approach 0 continuously, but, the order parameter C increases from 0 to a finite value discontinuously, as shown in Fig. 2(b) and (d). The softening phonon mode is a manifestation of a lattice instability just like structural phase transitions. A stability is restored by a condensation of the unstable mode. It results in a non-zero expectation value of the phonon operator ($\langle a \rangle \neq 0$), which may be taken as an order parameter for the M-BPI transition. The phonon hardens back to the bare mode as g is increased above g_{c2} because the screening is not effective in an insulating state. The a and $(n-1)$ are linearly coupled in the Hamiltonian of Eq. (1), and a zero/nonzero $\langle a \rangle$ is expected to lead to a zero/nonzero C . Therefore, an equally good order parameter is the C of Eq. (3).

In Fig. 2(b), we show z , C , and $\frac{\omega_0}{g}\langle a \rangle$ with increasing g . Note that $\langle n - 1 \rangle \approx \frac{\omega_0}{g}\langle a \rangle$ as expected from a simple mean-field theory. In Fig. 2(c), we plot the renormalized electron spectral function, $A(\omega) = -\frac{1}{\pi} \text{Im} \langle c, c^\dagger \rangle_\omega$, to show the MIT as g is increased along the cut 2b. In Fig. 2(d), we plot the renormalized phonon spectral function, $B(\omega) = -\frac{1}{\pi} \text{Im} \langle a + a^\dagger, a + a^\dagger \rangle_\omega$, calculated as g is increased using the method in [10]. Note that at $g = 0.1664$ which is close to $g_{c2} = 0.1665$, the phonon mode splits into two components as reported by Jeon *et al.* for the Anderson-Holstein model: one component develops into the soft mode and the other hardens back to the bare mode [10]. But, in the coexistence region where g_{c1} and g_{c2} are not close, the 1st order MIT from M to BPI pre-empts the emergence of the soft mode as g is increased because the MIT occurs at g_c ($g_{c1} < g_c < g_{c2}$).

The M-BPI transition may be understood from the Ginzburg-Landau theory. The free energy may be writ-




FIG. 3: In Fig. (a), we plot z as U is increased, and $\langle n_{\uparrow} n_{\downarrow} \rangle$ as U is increased and decreased along the cut 3 of Fig. 1. In Fig. (b), the electron spectral functions $A(\omega)$ are shown as U is increased along the same cut.

ten in terms of the the local phonon coordinate $\xi = \sqrt{\frac{\hbar}{2M\omega_0}}(a^\dagger + a)$ as

$$F = \frac{1}{2}\alpha\xi^2 + \frac{1}{4}\beta\xi^4 + \frac{1}{6}\gamma\xi^6, \quad (4)$$

with $\gamma > 0$. The phase transition can be 1st order for $\beta < 0$. In the context of the M-BPI transition, the $g_{c1} < g < g_{c2}$ regime corresponds to $\beta^2/4\gamma > \alpha > 0$, for which three local minima exit as a function of ξ . The g_{c1} , g_c , and g_{c2} correspond, respectively, to $\alpha = \beta^2/4\gamma$, $3\beta^2/16\gamma$, and 0. Although the α and β are complicated functions of U and g , the DMFT phase diagram of Fig. 1 indicates that α and β are mainly determined by, respectively, g and U . This observation suggests an interesting possibility that the M-BPI transition may become 2nd order for small values of U where $\beta \geq 0$, and a quantum tricritical point exists between E and T. This possibility will be explored further in a subsequent study.

(iii) *Transition between M and MHI.* – As the MIT from M to BPI is driven by growing charge susceptibility, the one from M to MHI is driven by growing spin susceptibility. In Fig. 3(a), we plot z as U is increased, and $\langle n_{\uparrow} n_{\downarrow} \rangle$ as U is increased and decreased for a fixed $g = 0.1$. z goes to 0 continuously, and $\langle n_{\uparrow} n_{\downarrow} \rangle_M = \langle n_{\uparrow} n_{\downarrow} \rangle_{MHI}$ as $U \rightarrow U_{c2}$. The M-MHI tran-




FIG. 4: The order parameters C is plotted as a function of g for the fixed $U = 2.5$ corresponding to the cut 4 of Fig. 1. The phase transition between BPI and MHI occurs along $U = E_{bp}$.

sition at $U_c = U_{c2}$ is, therefore, 2nd order. The phonons are hardly renormalized (not shown) because the charge degree of freedom is not soft unlike the MIT between M and BPI. In Fig. 3(b), $A(\omega)$ is plotted to show the MIT as U is increased.

The U_c does depend on g , although weakly, as can be seen from the Fig. 1, which implies that the M-MHI transition is coupled with the phonons. This raises an interesting possibility that some discrepancies between the experimental observations on V_2O_3 systems and DMFT results of the Hubbard model may be resolved in terms of the electron-phonon coupling [14].

(iv) *Transition between BPI and MHI.* – Now, we turn to an interesting question about the two kinds of insulators: Are BPI and MHI different, or, put differently, can the two insulating states be adiabatically connected? Along the line $U = E_{bp}$, the effective interaction in the static limit vanishes. One might then naively expect that a metallic state is the ground state along this line all the way to $U = E_{bp} \rightarrow \infty$. One can show, however, that for $U = E_{bp} \rightarrow \infty$ the ground state is an insulating state using the exact atomic limit Green's function [10, 15], as found from the detailed NGR+DMFT calculations.

In Fig. 4, we show C as g is varied for $U = 2.5$. The C changes discontinuously exactly at $E_{bp} = U$ and does not exhibit any other non-analytic behavior as g is varied. We checked another point on the line $E_{bp} = U$ by varying U with a fixed $g = 0.3$. The transition occurs exactly at $E_{bp} = U$ (not shown). This confirms that the two insulating states are distinct, and the phase transition between them occurs along the line of $U = E_{bp}$ and is 1st order. The point T in Fig. 1 is a triple point where two 1st order and one 2nd order phase transition lines meet.

We would like to thank Tae-Suk Kim, Yunkyu Bang, Ralf Bulla, Alex Hewson, and Dietrich Meyer for helpful comments and discussions. This work was supported by the Korea Science & Engineering Foundation (KOSEF)

through grant No. R01-1999-000-00031-0 and Center for Nanotubes and Nanoscale Composites (CNNC), and by the Ministry of Education through BK21 SNU-SKKU Physics program. JHH and HCL acknowledge the support from Center for Strongly Correlated Materials Research (CSCMR).

-
- [1] M. Imada, A. Fujimori, Y. Tokura, Rev. Mod. Phys. **70**, 1039 (1998).
 - [2] W. Metzner, D. Vollhardt, Phys. Rev. Lett. **62**, 324 (1989); *ibid*, 1066 (1989).
 - [3] A. Georges, G. Kotliar, W. Krauth, M. J. Rozenberg, Rev. Mod. Phys. **68**, 13 (1996).
 - [4] R. Bulla, Phys. Rev. Lett. **83**, 136 (1999).
 - [5] H. R. Krishna-murthy, J. W. Wilkins, K. G. Wilson, Phys. Rev. B **21**, 1003 (1980).
 - [6] K. G. Wilson, Rev. Mod. Phys. **47**, 773 (1975).
 - [7] R. Bulla, T. A. Costi, D. Vollhardt, Phys. Rev. B **64**, 045103 (2001).
 - [8] D. Meyer, A. C. Hewson, R. Bulla, Phys. Rev. Lett. **89**, 196401 (2002).
 - [9] R. Bulla, A. C. Hewson, Th. Pruschke, J. Phys.: Condens. Matter **10**, 8365 (1998).
 - [10] G. S. Jeon, T. H. Park, H. Y. Choi, Phys. Rev. B **68**, 045106 (2003).
 - [11] M. Capone, S. Ciuchi, Phys. Rev. Lett. **91**, 186405 (2003).
 - [12] G. Moeller, Q. Si, G. Kotliar, M. Rozenberg, D. S. Fisher, Phys. Rev. Lett. **74**, 2082 (1995).
 - [13] P. Benedetti, R. Zeyher, Phys. Rev. B **58**, 14320 (1998).
 - [14] P. Limelette *et al.*, Science **302**, 89 (2003).
 - [15] A. C. Hewson, D. Meyer, J. Phys.: Condens. Matter **14**, 427 (2002).